



The Rajasthan Medical Journal

Indexed in Index Copernicus -(ICV- 2020- 50.77)
Official Publication of the Medical and Health Department, Government of Rajasthan
Published by S.M.S. Medical College, Jaipur

Volume : 16

ISSN : 0485-9561

Issue 2 : April 2022

ORIGINALARTICLE

- Thymoma Associated Paraneoplastic Encephalitis: A Potentially Fatal but Treatable Entity

REVIEWARTICLE

- Monkeypox Disease and Vaccine: A Comprehensive

CASE REPORT

- Off Pump Total Arterial Coronary Revascularization in Dextrocardia with Situs InversusTotalis: A Case Report
- An Unusual Case Of Abdominal Wall Hematoma After Renal Allograft Biopsy
- Lesch Nyhan Syndrome



The Rajasthan Medical Journal



Editor in-chief

Dr. Sudhir Bhandari

Editors

Dr. S. M. Sharma
Dr. Deepak Mathur
Dr. Amarjeet Mehta

Co-Editors

Dr. Rajesh Sharma
Dr. Vinay Malhotra
Dr. I. D. Gupta
Dr. R. B. Sharma
Dr. S. Sitaraman
Dr. Amitabh Dube
Dr. Monica Jain

Associate Editors

Dr. Anita Singhal
Dr. Kapil Gupta
Dr. Jitendra Kumar Gupta

Assistant Editors

Dr. Sunil Mahavar
Dr. Laxmi Kant Goyal
Dr. Shruti Bhargava
Dr. Deepali Pathak
Dr. Maneesha Bhargava
Dr. Dhanraj Bagri
Dr. Puneet Agarwal

Advisory Editorial Board

Dr. R. K. Janav
Dr. Nupur Hooja
Dr. Ranjana Solanki
Dr. Sudhir Mehta
Dr. Raman Sharma
Dr. Vandana Mangal
Dr. Narendra Joshi
Dr. Praveen Mathur
Dr. Sandhya Gulati
Dr. Shivam Priyadarshi
Dr. M. L. Gupta
Dr. Pawan Singhal
Dr. Rakhi Arya

Specialty Sub- Committee

Dr. Reema Meena
Dr. Usha Jaipal
Dr. Sandhya Mishra
Dr. Seema Mehta
Dr. Bharti Malhotra
Dr. Lalit Batra
Dr. Vijay Pathak
Dr. Sandeep Nijhawan
Dr. Puneet Bhargava
Dr. Rashim Kataria
Dr. Mohnish Grover

Editorial Office

Medical Education Unit, S.M.S. Medical College, J.L.N. Marg, Jaipur – 302004

91-141-2518535. e-mail : medicaljournal.rajasthan@gmail.com

Editorial Assistants

Manish Jain, Vinod Kumar Sharma

Published by S.M.S. Medical College, Jaipur, Rajasthan, printed at Government Press, Jaipur Rajasthan

While every effort is made by the editorial team, members of boards and the publishers to avoid inaccurate or misleading information appearing in the individual articles and additional, the responsibility for the same solely rests with the authors concerned. The RMJ, the publishers, and members of the editorial team and board accept no liability whatsoever, for the consequences of any such inaccurate or misleading data, opinion or statement.



The Rajasthan Medical Journal

Volume : 16

ISSN : 0485-9561

Issue 2 : April 2022

	<i>Contents</i>	<i>Page No.</i>
ORIGINAL ARTICLE	Thymoma Associated Paraneoplastic Encephalitis: A Potentially Fatal but Treatable Entity.....	01
REVIEW ARTICLE	Monkeypox Disease and Vaccine: A Comprehensive	05
CASE REPORT	Off Pump Total Arterial Coronary Revascularization in Dextrocardia with Situs InversusTotalis: A Case Report.....	10
	An Unusual Case Of Abdominal Wall Hematoma After Renal Allograft Biopsy.....	16
	Lesch Nyhan Syndrome.....	18
The RMJ Policy & Guidelines	Instructions for Authors Renewal.....	21

ORIGINAL ARTICLE

Thymoma Associated Paraneoplastic Encephalitis: A Potentially Fatal but Treatable Entity

B.L. Kumawat*, Ruchi**, Piyush Savaliya**, Mridula Singh**

ABSTRACT

A 20 year old male presented with headache and seizure episodes in SMS Medical College, Jaipur. His chest X-ray and computed tomography revealed a right anterior mediastinal mass. MRI brain revealed T2/FLAIR cortical and subcortical hyperintensities. He underwent Thymectomy and was diagnosed with Type B2 thymoma histologically. The condition of patient improved and he recovered completely and became fully oriented after Thymectomy. Thymoma associated paraneoplastic encephalitis (TAPE) is a rare disease entity having very few case reported worldwide. Hereby, we are reporting a case of paraneoplastic encephalitis associated with thymoma.

Key words: Thymoma, paraneoplastic encephalitis, thymectomy

INTRODUCTION

Paraneoplastic neurological syndromes (PNS) are rare disorders associated with tumor, not caused by direct invasion, metastasis or consequences of treatment. These are found in <1% of patients with a malignancy. The neurological symptoms precede the tumor diagnosis in approximately 65% of patients with PNS. 50% of patients with thymoma have paraneoplastic neurologic syndromes myasthenia gravis being the most common. There are very few case reports of thymoma-associated paraneoplastic limbic or extralimbic encephalitis that can cause progressive neurologic decline and significant mortality without treatment. We report a case of a thymoma presenting as paraneoplastic encephalitis in a 20 year old male.

Clinical Summary

*Senior Professor, **Resident

Department of Neurology, SMS Medical College Hospital, Jaipur.

Corresponding Author

Dr Ruchi, DM Resident, Department of Neurology, SMS Medical College Hospital, Jaipur.

Email : dr.ruchijagota@gmail.com

Phone : 9996197957

A previously asymptomatic 20 year old male presented with 20 days history of headache followed by 1 day history of 3 episodes of generalized tonic clonic seizure and altered sensorium. Brain magnetic resonance imaging (MRI) showed bilateral foci of cortical and subcortical nonenhancing signal abnormalities on T2-weighted images (Figure 2). Chest x ray showed mediastinal widening with right hilar mass. Chest computed tomography (CT) revealed a large anterior mediastinal mass (Figure 1) that, on biopsy, showed type B2 thymoma (figure 3). Cerebrospinal fluid detected a predominantly lymphocytic pleocytosis. CSF was planned for antineuronal autoantibodies in the setting of a Thymic neoplasm but could not be done due to financial constraints of family.

Patient was managed with iv antiepileptics, steroids (IV Methyl prednisolone pulse) and IVIG (2gm/kg total dose). However, condition of patient continued to deteriorate, so, emergency Thymectomy was performed after obtaining high risk consent from the family. He started recovering after around 5 post operative days and became fully conscious and oriented at 10th post op day. Repeat MRI brain done two months later showed complete resolution of all lesions (Figure 4). Two years after the operation, there is no evidence of recurrence of tumor (Figure 5) and patient is seizure free presently.

DISCUSSION




Figure 1 showing MRI brain with T2 hyperintensities involving both cerebral hemispheres and cerebellum with no contrast enhancement.




Figure 2 showing preoperative Chest Xray and CT Chest showing anterior mediastinal mass.




Figure 3 showing histopathology and IHC(pancytokeratin and Tdt positive epithelial cells) suggestive of thymoma




Figure 4 showing resolution of lesions on MRI Brain done 2 months after Thymectomy.




Figure 5 showing post operative Chest Xray and CT chest of the patient after 2 years

Thymoma is the most common type of tumor in the anterior mediastinum. The usual age of presentation is 40- 60 years of age (M=F). Myasthenia gravis is the most common paraneoplastic syndrome associated with thymoma¹. First case of paraneoplastic encephalitis in the setting of thymic cancer was reported in 1988. Since then, very few case reports of thymoma associated paraneoplastic encephalitis (TAPE) have been reported² with no case reported from India. In most patients, neurologic manifestations include memory loss, confusion, and seizures. Seventy percent exhibited bilateral foci of nonenhancing hyperintensity on brain MRI, some appearing weeks after onset of symptoms. The differentials include metastases, primary brain tumors, and infection². The presence of a mediastinal mass and pathologic confirmation of thymoma suggests a paraneoplastic syndrome in our case. Patients with Thymoma-associated encephalitis most commonly have VGKC-complex Abs-LGI1 and Caspr2, with more likelihood of Caspr2- positivity³ although other antibodies to CRMP-5, anti GAD and GABA-A Receptor have also been described in literature^{4,5}. The diagnosis of TAPE is based on the combination of the Encephalitis syndrome, MRI findings, and detection of antineuronal autoantibodies in the setting of a Thymic neoplasm². The limitation of this case is inability to get CSF antineuronal antibodies done owing to non affordability of the family for the same.

Immediate intervention is necessary to manage TAPE in order to prevent progressive neurologic decline and death. Immunotherapy with IgG and corticosteroids will improve neurologic symptoms resulting from inflammatory response, but complete resection of the thymoma confers the best chance at disease-free survival from oncologic as well as paraneoplastic standpoints⁶.

Plasmapheresis can be done in refractory cases⁷.

It may take weeks to months after Thymectomy for symptom resolution⁸. Outcome is driven more by ongoing neurologic insult than the underlying thymic disease. The mortality rate in TAPE is thirty-three percent, due to the nervous system compromise⁹. Only 21% of patients with TAPE achieve symptom-free survival; all of these patients had complete resection of their thymoma².

It is necessary on the part of clinician to make an early link between thymoma and encephalitis which can lead on to immediate intervention in the form of complete thymoma resection giving patient the best chance of disease free survival.

Acknowledgments- Late Dr CM Sharma- for his exceptional guidance and support,

Dr Raj Kumar Yadav- Senior Professor, CTVS deptt, SMS Medical College and Hospital, Jaipur- for his efficient management

REFERENCES

1. Rizzardi G, Campione A, Scanagatta P, Terzi A. *Paraneoplastic extra limbic encephalitis associated with thymoma*. Interact Cardiovasc Thorac Surg 2009;9:755-6.
2. Erkmen CP, Camilo E, Fadul CE, Dalmau J, and Erkmen K et al- Thymoma-associated paraneoplastic encephalitis (TAPE): Diagnosis and treatment of a potentially fatal condition- J Thorac Cardiovasc Surg 2011;141:e17-20
3. Lorio R, Spagni G, Evoli A-Paraneoplastic neurological syndromes associated with mediastinal tumors - *Mediastinum* 2018;2:8
4. Yu Z, Kryzer TJ, Griesmann GE, Kim K, Benarroch EE, Lennon VA. CRMP-5 neuronal antibody: marker of lung cancer and thymoma-related autoimmunity. Ann Neurol. 2001; 49:146-54.
5. Petit-Pedrol M, Armangue T, Peng X, Bataller L, Cellucci

- T, Davis R, McCracken L, Martinez-Hernandez E, Mason WP, Kruer MC, Ritacco DG, Grisold W, Meaney BF, Alcala C, Sillevs-Smitt P, Titulaer MJ, Balice-Gordon R, Graus F, Dalmau J. Encephalitis with refractory seizures, status epilepticus, and antibodies to the GABAA receptor: a case series, characterisation of the antigen, and analysis of the effects of the antibodies. *Lancet Neurol.* 2014; 13:276–86.
6. Antoine JC, Honnorat J, Anterior CT, Aguera M, Absi L, Fournel P, et al. Limbic encephalitis and immunological perturbations in two patients with thymoma. *J Neurol Neurosurg Psychiatr.* 1995;58:706-10.
7. Rickman O, Parisi J, Yu Z, et al. Fulminant autoimmune cortical encephalitis associated with thymoma treated with plasma exchange. *Mayo Clin Proc* 2000;75:1321–6.
8. Suh JW, Haam SJ, Song SW, Paraneoplastic Encephalitis Associated with Thymoma: A Case Report. *Korean J Thorac Cardiovasc Surg* 2013;46:234-6.
9. Hartmann CA, Roth C, Minck C, Niedobitek G. Thymic carcinoma. Report of five cases and review of the literature. *J Cancer Res Clin Oncol.* 1990;116:69-82.

REVIEW ARTICLE

Monkeypox Disease and Vaccine: A Comprehensive

Monica Jain*, Anil Bhandari**, Abhinav Agrawal***

ABSTRACT

While the world was still recovering from horrors of the Coronavirus disease 2019 (COVID-19), another viral disease, this time the Monkeypox virus, a member of Orthopoxvirus genus¹ of the Poxviridae family started spreading. Monkeypox is a zoonotic infection with rare instances of human-to-human transmission reported worldwide². Monkeypox gets its name from the Macaque monkeys in whom this viral infection was first witnessed³. Owing to an outbreak in 2022, Monkeypox virus disease which was previously known to be endemic to certain regions of Africa was reported from regions of Europe and western hemisphere⁴. Extensive research and advent of vaccines have been the front-runners in containing the outbreak.

INTRODUCTION

Monkeypox disease is an infectious zoonotic disease caused by Monkeypox virus, a member of Poxviridae family^{1,2}. It belongs to genus Orthopoxvirus, same as that of Smallpox, Cowpox and Vaccinia virus⁵. Like the other members of Poxviridae family, Monkeypox viruses also are large, enveloped, linear double stranded DNA viruses with a genome size ranging from 130-360 kbp⁶. Several animal species are known to be susceptible to the Monkeypox virus like squirrels, Gambian pouched rats, dormice, and non-human primates².

Monkeypox virus was first observed in laboratory monkeys in Copenhagen, Denmark in 1958⁷. However the first human case was not reported till 1970 when a 9-month-old boy was found positive for the virus during mass screening for the Smallpox virus in Zaire (now known as Democratic Republic of Congo)⁸.

The first case of the recent outbreak of Monkeypox disease was reported on May 6, 2022 in

United Kingdom. The outbreak spread out from Central and West Africa to other continents⁹. On July 23, 2022, WHO declared the outbreak a public health emergency of international concern (PHEIC)¹⁰. As of September 30, 2022, 12 cases of Monkeypox disease have been reported in India in addition to 68,428 confirmed cases in 106 countries worldwide⁹.




Fig 1: Worldwide distribution of Monkeypox

Photo Credit: /www.cdc.gov/poxvirus/monkeypox/response/2022/world-map

Epidemiology:

Monkeypox disease is originally known to be endemic to Sub-Saharan Africa for hundreds of years with thousands of cases reported annually. The disease became a matter of global concern for public health in 2003, when an outbreak was reported in United States of America linked to infective pet dogs¹¹. The natural host and the reservoir of the Monkeypox virus still remains a mystery with studies suggesting both humans and monkeys as incidental hosts.

The Monkeypox virus is known to have two clades: West Africa (WA) clade and the Congo Basin (CB) clade with the latter known to be more virulent. However

* Senior Professor, **PG Student, ***3rd Year MBBS
Department of Pharmacology, SMS Medical College, Jaipur

Corresponding Author

Dr. Anil Bhandari

PG Student, Department of Pharmacology, SMS Medical College, Jaipur

Contact No. 9717545233 Email : anil.bhandari@gmail.com

in the current outbreak, West African clade seems to be responsible for the sudden rise of cases in Europe and other parts of the world¹². Epidemiological studies suggest an R_0 value (degree of transmissibility of disease) ranging from 1.10 to 2.40 for the disease with case fatality rate estimated at 0-3% for West African clade and 0-11% for Congo Basin clade¹¹.




Fig 1: Case distribution of the two clades over the years.

Photo Credit: **The changing epidemiology of human monkeypox—A potential threat? A systematic review**
Transmission and Pathogenesis:

Monkeypox virus primarily infects invertebrates and lower mammals. Animal-to-human transmission is the chief reason for presence of disease in humans. Animal-to-human transmission can result either from direct contact with blood and body fluids or from cutaneous and mucosal lesions of infected animals¹³. Human-to-human transmission can occur from close contact with respiratory secretions, or skin lesions of an infected person. Transplacental transmission from mother to fetus is also a possibility and may give rise to Congenital Monkeypox disease. Few cases of perinatal transmission have also been reported. The sexual route of transmission is still under evaluation with early reports from certain studies suggesting higher prevalence in men who have sexual contact with other men¹¹.

The incubation period for the disease ranges from 5-21 days with person being non-contagious throughout this period implying that infection is spread only after symptoms appear. The virus multiplies at the site of inoculation before entering the blood stream. Lesions appear 2-3 days after onset of fever and are mainly maculo-papular in nature. Serum antibodies also become positive by this period¹⁴.




Fig 2: Guide Map to Monkeypox disease

Photo Credit: Emergence of monkeypox: Risk assessment and containment measures ;**June 2022;**
O.P.Chaudhary et al.

Clinical Features:

Monkeypox is usually a self-limiting disease with symptoms lasting from 2 to 4 weeks. Severe cases occur more commonly among children and are related to the extent of virus exposure, patient health status and nature of complications. Underlying immune deficiencies like diabetes mellitus and AIDS may lead to more detrimental outcomes¹⁴.

The initial five days called invasion period manifests with fever, headache, lymphadenopathy, back pain, myalgia and asthenia. Lymphadenopathy is a typical feature of monkeypox distinguishing it from the early stages of similar diseases like chickenpox, measles and smallpox^{11,13}.

The skin eruption commonly appears within 1-3 days after the onset of the fever. The rash appears predominantly on the face and extremities sparing the trunk. Lesions can also be seen on the oral mucosa, genitalia, conjunctiva as well as the cornea¹⁴.

The rash evolves in textbook fashion from flat base to macules to papules to vesicles to pustules and crusts which in due course eventually dry up and fall off.



Fig 3: Examples of Monkeypox disease Rashes

Photo credit: UK Health Security Agency

Complications:

Complications of Monkeypox disease include secondary infections, bronchopneumonia, sepsis, encephalitis, and infection of the cornea with ensuing loss of vision. However the extent to which the asymptomatic infection may occur remains unknown¹⁵.

Lab Diagnosis and Differential Diagnosis:

Polymerase Chain Reaction (PCR) remains the diagnostic modality of choice for the disease. Samples are taken from active skin lesions. PCR is believed to have higher sensitivity even in presence of bacterial contamination¹¹. A hybridization assay, utilizing probe (MGB Eclipse trade mark probe), targets envelope protein gene (B6R) and specifically detects Monkeypox virus (MPXV)¹⁶.

Diseases like chickenpox, measles, bacterial skin infections, scabies, syphilis and medication-associated allergies which manifest with rashes serve as important differential diagnoses for the Monkeypox virus disease owing to the presence of classical rashes in the Monkeypox disease too. However the typical rash pattern and presence of lymphadenopathy in prodromal stages of the illness are the characteristic features of the Monkeypox disease¹⁴.

Treatment:

The management of Monkeypox disease is

primarily based on providing symptomatic relief and at the same time preventing potentially fatal complications. DNA Polymerase inhibitors like Cidofovir known to be effective in animals are under trial for human use in Monkeypox disease¹¹. Recently, a drug Tecovirimat which inhibits **p37**, a highly conserved protein in all orthopoxviruses known to mediate the formation of enveloped virions, has been approved for use against Monkeypox disease in various countries^{11,14,17}.

United States Food and Drug Administration has also licensed Vaccinia Immunoglobulins (known to be effective against Orthopoxviruses) for effective management and adequate prophylaxis against the fast spreading disease¹¹.

Vaccines and Vaccination:

Vaccines against Smallpox disease have been demonstrated to be about 85% effective in preventing Monkeypox disease. A still newer vaccine based on a modified, attenuated, Vaccinia virus strain (Ankara strain) was approved for the prevention of Monkeypox in 2019. This vaccine is a third generation smallpox virus and is given in two-doses for which availability remains limited^{18,19}. Live attenuated, non-replicating smallpox and Monkeypox vaccine that elicits both humoral and cellular immune response to orthopoxviruses is under clinical trials.

Vaccination is indicated for the prevention of Monkeypox disease in adults who are at high risk of contracting the infection. It has been 40 or more years since all countries stopped routine smallpox vaccination with vaccinia-based vaccines that protected against Monkeypox. The population unvaccinated against smallpox which is born after 1980 is more susceptible to Monkeypox virus infection.

Dosing:

- a) Subcutaneous Administration: 0.5 mL SC x2 doses; 4 weeks apart
- b) Intradermal Administration: 0.1 mL ID x2 doses; 4 weeks apart (Emergency Use Authorization)¹⁸

Adverse Effects:

Local side effects include injection site pain, redness, swelling, induration and itching. Systemic manifestations include muscle pain, headache, fatigue, nausea and chills.

Pregnancy:

Human data are insufficient to inform vaccine-associated risks in pregnancy. Studies in animals revealed no evidence of harm to the fetus.

Lactation:

Unknown if excreted in human milk; data are not available to assess vaccine effects in breastfed infants or on milk production/excretion^{18,19,20}.

Prevention:

Forewarned is forearmed therefore educating people about risk factors and mitigating the exposure to the virus to minimum are chief strategies for prevention. Due to reported human-to-human transmission by respiratory secretions, isolation of the diseased and Quarantine of those exposed also seems as a strong strategy.

The Ministry of Health and Family Welfare (MoHFW), Government of India has already released guidelines on the detection and management of Monkeypox disease. The National Institute of Virology (NIV) in Pune has been designated as nodal laboratory for Monkeypox virus testing in India¹¹.

REFERENCES

1. WHO(World health organization). Geneva. 2022. fact sheet Monkey Pox. [Updated 2022 Sep 9][cited 2022 Sep 9]www.who.int/news-room/fact-sheets/detail/monkeypox/May
2. CDC (centers for disease control and prevention). United States. Veterinarian monkeypox in animals. [updated 2022 Sep 9] [cited 2022 Sep 9] www.cdc.gov/poxvirus/monkeypox/veterinarian/monkeypox-in-animals.
3. Lindsay Smith Rogers, John Hopkins University. Maryland. 2022 what you need to know about monkey pox. [updated 2022 Sep 9] [cited 2022 Sep 9] https://publichealth.jhu.edu/2022/what-you-need-to-know-about-monkeypox/
4. WHO(World health organization). Geneva. 2022. Disease Outbreak News . [Updated 2022 Sep 9] [cited 2022 Sep 9] https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON396
5. Charles Patrick Davis, MD, PhD. www.medicinenet.com/orthopox/definition/[updated 2022 Sep 9]
6. National Institute of Health(NIH), US [Updated 2022 Sep 9] [cited 2022 Sep 9]https://ncbiinsights.ncbi.nlm.nih.gov/2022/05/26/monkeypox-virus-genome
7. Moore MJ, Rathish B, Zahra F. Monkeypox. 2022 Oct 8. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. PMID: 34662033.
8. John Thornhill, M. D. , Ph. D. , et al. (n.d.). Monkeypox Virus Infection in Humans across 16 Countries .April–June 2022
9. CDC (Centers for disease control and prevention). United States. Monkeypox response. [updated 2022 Sep 9] [cited 2022 Sep 9]cdc.gov/poxvirus/monkeypox/response/2022 Monkeypox Outbreak Global Map
10. Nuzzo JB, Borio LL, Gostin LO. The WHO Declaration of Monkeypox as a Global Public Health Emergency. *JAMA*. 2022;328(7):615–617. doi:10.1001/jama.2022.12513
11. Lahariya C, Thakur A, Dudeja N. Monkeypox Disease Outbreak (2022): Epidemiology, Challenges, and the Way Forward. *Indian Pediatr*. 2022 Aug 15;59(8):636–642. doi: 10.1007/s13312-022-2578-2. Epub 2022 Jun 27. PMID: 35762024; PMCID: PMC9419123.
12. World Health organization. Multi-country

- Monkeypox outbreak: Situation update. Accessed June 17, 2022. Available from: <https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON393>
13. Titanji, B. K., Tegomoh, B., Nematollahi, S., Konomos, M., & Kulkarni, P. A. (2022). Monkeypox: A Contemporary Review for Healthcare Professionals. *Open Forum Infectious Diseases*, 9(7), ofac310. <https://doi.org/10.1093/ofid/ofac310>
14. Kaler, J., Hussain, A., Flores, G., Kheiri, S., & Desrosiers, D. (2022). Monkeypox: A Comprehensive Review of Transmission, Pathogenesis, and Manifestation. *Cureus*, 14(7), e26531. <https://doi.org/10.7759/cureus.26531>
15. Dhawan M, Priyanka, Choudhary OP. Emergence of monkeypox: Risk assessment and containment measures. *Travel Med Infect Dis*. 2022 Sep-Oct; 49:102392. doi: 10.1016/j.tmaid.2022.102392. Epub 2022 Jun 22. PMID: 35752293; PMCID: PMC9217686.
16. Li, Y., Olson, V. A., Laue, T., Laker, M. T., & Damon, I. K. (2006). Detection of monkeypox virus with real-time PCR assays. *Journal of Clinical Virology: The Official Publication of the Pan American Society for Clinical Virology*, 36(3), 194–203. <https://doi.org/10.1016/j.jcv.2006.03.012>
17. John E. Bennett MD, in Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases, 2020, Miscellaneous Antiviral Agents (Interferons, Tecovirimat, Imiquimod, Poxapavir, Pleconaril) [cited 2022 Sep 9]
18. Townsend, M. B., Keckler, M. S., Patel, N., Davies, D. H., Felgner, P., Damon, I. K., & Karem, K. L. (2013). Humoral immunity to smallpox vaccines and monkeypox virus challenge: proteomic assessment and clinical correlations. *Journal of Virology*, 87(2), 900–911. <https://doi.org/10.1128/JVI.02089-12>
19. Medscape, [US], Jynneos-smallpox-vaccinia-monkeypox-vaccine-live-nonreplicating <https://reference.medscape.com/drug/jynneos-smallpox-vaccinia-monkeypox-vaccinelive-nonreplicating-4000029> [cited 2022 Sep 9]
20. Jynneos (smallpox and monkeypox vaccine, live, non-replicating) package insert. K4vistgaard, Denmark: Bavarian Nordic A/S; 2021 June.

CASE REPORT

Off Pump Total Arterial Coronary Revascularization in Dextrocardia with Situs Inversus Totalis: A Case Report

Hemlata Verma*, Anula Sisodia*, R M Mathur**, Himanshu Mahala*, Vaibhav Gupta***, Deva Ram Chaudhary***, Harnish Singh Bhatia***

ABSTRACT

Dextrocardia with situs inversus is a rare entity and if not associated with any congenital heart disorder then these persons lead a normal life. CABG in dextrocardia is technically a challenging task. Here we are reporting a case of off pump total arterial CABG in a patient with dextrocardia with situs inversus totalis with regard to technique and modifications for the surgery.

Key words: dextrocardia, situs inversus totalis, CABG, off pump, congenital, heart.

INTRODUCTION

Dextrocardia is usually associated with congenital heart diseases and if not, then, it behaves normally like levocardia with regard to natural history. Dextrocardia with situs inversus behaves more like normal than dextrocardia with situs solitus.

Coronary artery disease after age 40 year is very common in normal levocardia heart (i.e. apex of heart is on left side with atrial situs and main mass of heart is lying left of the midline) although may occur in younger age group. Dextrocardia is not exception for this yet an uncommon entity.

Dextrocardia is associated with surgical challenges eg, position of surgeon, cannulation for CPB, conduit selection, graft configuration, sequence of anastomosis etc.

Here, we are presenting a case of situs inversus totalis with dextrocardia with TVD in which total arterial revascularization was done off pump.

PATIENT PROFILE

59 year old female presented to us with history of recurrent angina right side of chest radiating to right arm for last 3 years and an episode of acute myocardial

infarction 3 months back. Coronary angiography was done which showed Tripple vessel disease. She was referred to us for surgery.

She was non hypertensive and, non diabetic with no history of other chronic illness.

Patient was thoroughly evaluated with routine blood investigations, viral markers, and covid test. Her chest x-ray (Picture/illustration 1) showed dextrocardia with situs inversus. Liver was under left hemidiaphragm and fundal shadow was under right hemidiaphragm.

HRCT thorax for covid infection was also negative

Electrocardiography (Figure 2) was typical of dextrocardia and Coronary artery disease i.e. inverted P wave with negative QRS deflection in lead I and upright P wave with positive QRS deflection in lead avR and poor progression of R wave in left chest leads.

Findings suggestive of Coronary artery disease were presence of q wave in lead V2, V5, V6, ST depression with T wave inversion in lead V4R, V5R, and V6R.

2D echo (Figure 3) showed situs inversus with dextrocardia with normal chamber dimensions. Superior vena cava and Inferior vena cava were on left side of spine and draining into left side placed Right atrium. Pulmonary veins draining into right sided left atrium. Apex was lying on right side of midline. Morphologic right ventricular was on left side of midline and morphologic left ventricle was on right side of midline. Aorta was lying on right side of spine and ascending aorta was situated on left side of main pulmonary artery with ventriculo arterial concordance. There was grade II left ventricular diastolic dysfunction with normal systolic function (Ejection fraction – 55 %).

*Associate Professor, **Senior Professor, ***Resident Doctor

Department of Cardiothoracic and Vascular Surgery, SMS Medical College & Hospital, Jaipur, Rajasthan

Corresponding author:

Dr. R M Mathur

Senior Professor and Unit Head

CTVS Department, SMS Hospital, Jaipur, Rajasthan

Address: 48, Bajaj Nagar Enclave, Jaipur, Rajasthan (302015)

+91-9829063353

Email : rajendra.mathur@rediffmail.com

Coronary angiography (Figure 4a,4b,4c,4d) revealed mirror image coronary artery pattern with left arterial system was lying on right side of main mass and right coronary artery was on left side of main mass. There was approximately 95% stenosis in proximal Left anterior descending artery, obtuse marginal 1, proximal right coronary and distal right coronary artery just before crux.

Ultrasonography abdomen confirmed the finding of abdominal situs that is liver was on left side and spleen was on right side.

There was moderate to severe obstructive respiratory dysfunction and both carotid arteries were normal on colour Doppler study.

With negative corona profile and HRCT thorax, patient was taken for surgery.

Her both radial arteries were checked with Allen test for harvesting IMA and Radial artery.

INTRA OPERATIVE PROFILE

ECG electrodes were fixed on back side of right chest wall. Patient was anesthetized with fentanyl, sodium thiopental, and rocuronium and maintained with sevoflurane and intermittent doses of fentanyl and rocuronium. Arterial and venous monitoring lines were placed and urinary catheterization was done.

Chest was opened with midline sternotomy, pericardium was opened and hitched up. Anatomy (Figure 5) of heart was consistent with mirror image pattern with right aortic arch. LAD was situated on right side, OM was on right lateral surface, RCA was on left side, and PDA was situated on inferior wall left side.

After inspection of anatomy, RIMA (Figure 6a,6b,6c), LIMA (Figure 6d,6e), and LEFT RADIAL (Figure 6f) artery were harvested on 1 mg/kg heparin dose with ACT more than 280 seconds. RIMA was used as pedicled graft and LIMA and RADIAL aretery were used as free graft. Patient was heparinized again to achieve ACT more than 300 seconds. Procedure was done off pump and with surgeon standing on right side of the patient. Proximal anastomosis (Figure 7) were done first with LIMA was anastomosed to right lateral surface and RADIAL artery was anastomosed to left anterolateral surface of ascending aorta with 7/0 prolene in continuous manner. Patency was checked.

Stabilizer was fixed on inferior arm of retractor and LAD (Figure 8) was stabilized. Exposure was improved with hitching up the pericardial stay taken in between the right superior and inferior pulmonary vein. A 5/0 prolene control sling was taken around LAD proximal

to the proposed site of anastomosis. Adequate sized LAD arteriotomy was done and shunt of 1.75mm size was placed inside the arteriotomy. RIMA was anastomosed to LAD as pedicled graft with 7/0 prolene in tension free manner. RIMA graft was fixed to epicardium. Now stabilizer was fixed on inferior arm of retractor near its right arm and OM1 artery (Figure 9) was fixed. LIMA was anastomosed to OM1 in similar manner as LAD grafting. For PDA exposure, octopus was stabilized at right arm of retractor near its inferior portion and RADIAL artery was anastomosed to PDA (Figure 10) similarly. Shunt size for OM1 and PDA was 1.5mm.

After completion of the procedure heparin was reversed. Hemodynamically patient's parameters were remain stable throughout the procedure therefore no inotropes were required. Chest was closed in usual manner after inserting bilateral pleural and mediastinal drains. She was shifted to ICU on nitroglycerine infusion.

Postoperative course was uneventful and she was extubated in 6 hrs. Patient was discharged on day 7.

DISCUSSION

Dextrocardia with situs inverses is an unusual congenital abnormality with an incidence of 1: 12,000¹.

It is subjected to coronary artery disease as in normal population with levocardia and situs solitus².

Literature shows that dextrocardia was first reported by anatomist surgeon Hieronymus Fabricius in 1606³, and Marco Aurelia Severinus in 1643 reported dextrocardia with situs inversus⁴. In adults, mostly it is dextrocardia with atrial and abdominal situs inversus and this is rarely associated with congenital cardiac anomalies.

CABG was first done in dextrocardia in 1980⁵.

Performing surgery when standing on right side of patient in dextrocardia is technically difficult and doing it off pump is like a pimple has grown upon an ulcer. Some issues have to be sorted out during surgery like where should surgeon stand, doing it off pump or on pump, cannulation strategies, conduits for LAD, OM, PDA, which coronary artery is to be anastomosed first or sequence of anastomosis, position of stabilizer etc.

Selection of conduit and configuration of graft is very important in CABG and it becomes very crucial in dextrocardia because now main mass of heart is on right side. RIMA can easily reach upto LAD in this situation; however Kuwata⁶ and chakravarthy⁷ used skeletonised LIMA in situ for LAD.

In most of the case reports and paper presented so far it was shown that one arterial graft, which is usually RIMA for LAD was used and for the rest of coronaries venous grafts were used. Kuwata T et al reported a case where he performed total arterial surgery off pump⁶. We decided to do a total arterial revascularization off pump. RIMA was used as pedicled graft for LAD, and LIMA and RADIAL artery conduits were used as free grafts for OM and PDA. Pancelet did total arterial revascularization with the help of cardiopulmonary bypass⁸.

Usually RIMA is anastomosed to LAD, considering its proximity to rightward LAD.

In dextrocardia, obtuse marginal vessels are more anterior and a conduit with short length can be used, therefore it is more suitable to do total arterial revascularization as they are less prone to kinking than SVGs⁹.

Because of position of the heart in dextrocardia it is preferred to stand on left side of the patient for the coronary anastomosis as it is easy to graft LAD and OM exposed by retracting the heart to left. Few cases are reported in which surgery was done when standing on right side¹⁰ and we found that this surgery can be done with standing on right side without any difficulty.

Texas study showed that in dextrocardia, most surgeons were on the left side, cardiopulmonary bypass was used frequently, and usually one arterial and rest were venous grafts were used¹¹.

When using cardiopulmonary bypass position of aortic arch is very important to place the arterial cannula and usually in dextrocardia it is right aortic arch¹². Presence of SVC draining into coronary sinus has to be checked for venous cannulation and retrograde cardioplegia¹³.

SUMMARY

Dextrocardia poses challenges to surgeons but slight modifications make the grafting easier. Although in our case we did not change the surgeon's position and completed the whole procedure as we do CABG in levocardia without any difficulty. By doing total arterial revascularization without the help of cardiopulmonary bypass we could do the best for the patient in our scenario.

REFERENCES

1. Bohun CM, Potts JE, Casey BM, et al. A population – based study of cardiac malformations and outcomes associated with dextrocardia. *Am J Cardiol* 2007;2013:305-9.
2. Hynes HM, Gau GT, Titus JL. Coronary heart disease in situs inversustotalis. *Am J Cardiol*. 1973;31:666 – 9.
3. Cleveland M. Situs inversusviscerum : Anatomic study. *Arch Surg*. 1926;13:343.
4. Brown JW. Congenital Heart Disease. London: *Staples Press Ltd*;1950.
5. Irvin RG, Ballenger JF. Coronary artery bypasses surgery in a patient with situs inversus. *Chest*. 1982;81:380- 1.
6. Kuwata T, Kawata T, Ueda T, Sakaguchi H, Nagasaka S, Taniguchi S. Off pump quintuple coronary artery bypass grafting for situs inversustotalis. *Jpn J ThoracCardiovascSurg* 2004;52(10):473-5.
7. Chakravarthy M, Jawali V, Nijagal D. Off- pump coronary artery bypass surgery in dextrocardia : a report of two cases. *Ann ThoracCardiovascSurg* 2008;14(3):187- 91.
8. Pancelet AJ, Dion R, Lengele B, Nairhomme P. Complete arterial revascularization in coronary artery bypass grafting in patients with situs inversustotalis. *J CardiovascSurg (Torino)* 2006;47(4):477 – 9.
9. Naik MJ, Chua YL, Ding ZP, Lau KW. Coronary artery bypass grafts in situs inversus. *J CardiovascSurg (Torino)* 2002;43(2):181 – 4.
10. Saad RA, Badr A, Goodwin AT, Dunning J. Should you stand on the left or the right of a patient with dextrocardia who needs coronary surgery? *Interact Cardiovasc Thoracic Surg*. 2009;9:698-702.
11. Murtuza B, Gupta P, Goli G, Lall KS. Coronary revascularization in adults with dextrocardia. Surgical implications of anatomic variants. *Tex Heart Inst J*. 2010; 37:633-40.
12. Arcilla RA, Gasul BM. Congenital dextrocardia. Clinical, angiographic, and autopsy studies on 50 patients. *JPediatr*. 1961;58:251-62.
13. Kurisu K, Ochiai Y, Kumeda H, Kimura S, Tanaka K, Tominaga R. Coronary artery bypass grafting with absent right superior vena cava in visceratrial situs solitus. *Jpn J ThoracCardiovascSurg* 2004;52(4):191-3.

[illegible][illegible]

The RMJ : Volume - 16, Issue- 2, April 2022




Figure 4c: LAD and Left Circumflex Artery




Figure 4d: Right Coronary Artery



Figure 5: Anatomy



Figure 6a: RIMA Harvesting



Figure 6b: RIMA Graft




Figure 6c: RIMA Graft




Figure 6d: LIMA Harvesting



Figure 6e: LIMA



Figure 6f: Left Radial Artery Graft Harvesting




Figure 7: Proximal Anastomosis




Figure 8: LAD Grafting



Figure 9: OM 1 Grafting



Figure 10: PDA Grafting

CASE REPORT

An Unusual Case Of Abdominal Wall Hematoma After Renal Allograft Biopsy

Geeta*, Rakesh Gupta**, Tushar Gupta*, Pankaj Beniwal***, Vinay Malhotra****, Dhananjai Agarwal****, SitaRam*, Niranjan Gogoi*

ABSTRACT

Renal biopsy is often required to diagnose allograft dysfunction. Although it's a safe procedure, complications do occur. We report an unusual case of anterior abdominal wall hematoma after renal allograft biopsy. The patient developed progressive swelling over allograft site. Initially patient was managed conservatively followed by percutaneous drain. Color doppler evaluation of overlying abdominal wall can help to look for significant vessels before biopsy procedure and avoid such complications.

Key Words: Renal biopsy, Allograft, Hematoma.

INTRODUCTION

Renal allograft biopsies are done for evaluation of allograft dysfunction after ruling out pre-renal and post renal causes. The use of ultrasound guidance has reduced the chances of significant bleeding. This is a case report of enlarging abdominal wall hematoma after percutaneous renal allograft biopsy which was managed with an abdominal wall drain.

CASE REPORT

A 46 year old female who underwent renal transplantation 12 years ago presented to our hospital with asymptomatic increase in serum creatinine. Systemic examination was unremarkable. Tacrolimus levels were normal. Renal biopsy was done, after ruling out pre-renal and post renal causes, for evaluation of graft dysfunction.

Renal allograft biopsy was done under USG guidance from the upper pole using an 18 G needle after checking the coagulation parameters. A tender swelling developed at the biopsy site 4-6 hours after the procedure which was gradually increasing in size. There was no associated hematuria. USG showed organised collection of 139x70 mm around transplanted kidney, likely

hematoma. Doppler study didn't show any evidence of pseudoaneurysm or arteriovenous fistula. Patient's hematocrit dropped and hence blood transfusions were given. The hematoma further increased in size the next day and CT abdomen was done which was suggestive of large hypodense collection in lateral abdominal wall (fig 1,2). CT angiography was done to look for the site of bleed. The bleeding vessel could not be identified.



Fig 1




Fig 2

*Resident, **Associate Professor, ***Professor, ****Senior Professor
Department of Nephrology, SMS Medical College & Hospital, Jaipur

Corresponding Author:

Dr. Shubham Agarwal

Resident Doctor

Department of Medicine, Mahatma Gandhi Medical College
Jaipur.

Email: drdhananjainephro@yahoo.com

Ph. No. 9414459790

The patient was managed conservatively with bed rest and regular monitoring of hematocrit. Hematoma stopped expanding 3 days later but due to persistent tenderness abdominal wall drain was inserted for early drainage of hematoma. Patient responded well to this management and was kept on regular follow up.

DISCUSSION

Renal biopsy is usually recognized as a safe procedure in native and transplant kidney.^{1,2} Biopsy procedures of renal allograft are known to have a higher incidence of gross hematuria, hematoma formation, hemoperitoneum and AV fistulae.^{3,4} While minor complications usually resolve spontaneously, major complications require further treatment, sometimes even evacuation of hematoma or nephrectomy.⁵

Injury to arteries in abdominal wall can occur during paracentesis, chest tube insertion and rarely following percutaneous biopsy.^{6,7,8} The arteries most commonly injured are the branches of superior epigastric artery, inferior epigastric artery or circumflex iliac artery.^{7,8,9} In this case we were suspecting an injury to anterior abdominal wall vessel which was responsible for the formation of hematoma although the exact site of bleed could not be identified. Knowledge of vasculature of anterior abdominal wall is useful in such circumstances and Doppler study can help to identify the source of bleed.

CONCLUSION

Abdominal wall hematoma is a rare complication of allograft biopsy. Although there have been quite a few case reports of renal and perirenal hematoma after allograft biopsy, anterior abdominal wall hematoma has been reported scarcely. The person performing biopsy should be aware that injury to abdominal wall vessels can occur during the procedure and these should be evaluated as the cause of bleed if no renal parenchymal lesion is obvious. A color Doppler evaluation of the abdominal wall overlying the allograft can be useful to diagnose such cases.

REFERENCES

1. Tondel C, Vikse BE, Bostad L, Svarstad E. Safety and complications of percutaneous kidney biopsies 1988-2010. *Clin J Am Soc Nephrol*. 2012;7:1591-1597.
2. Esposito V, Mazzon G, Baiardi P, et al. Safety and adequacy of percutaneous kidney biopsy performed by nephrology trainees. *BMC Nephrology*. 2018;19:14.
3. Parrish AE. Complications of percutaneous biopsy: a review of 37 years' experience. *Clin Nephrol*. 1992;38:135-41.
4. Gainza FJ, Minguela I, Lopez-Vida I, Ruiz LM, Lampreabe I. Evaluation of complications due to percutaneous renal biopsy in allografts and native kidneys with color-coded doppler sonography. *Clin Nephrol*. 1995;43:303-8.
5. Burstein DM, Korbet SM, Schwartz MM. The use of the automatic core biopsy system in percutaneous renal biopsies: a comparative study. *Am J Kidney Dis*. 1993;22:545-52.
6. Sadat U, Jah A, Ward N, Gaunt M. Superior epigastric artery pseudoaneurysm-A rare complication of chest drain insertion in coronary artery bypass grafting. *J Cardiothorac Surg* 2007;2:21.
7. Lam EY, McLafferty RB, Taylor LM Jr., Moneta GL, Edwards JM, Barton RE et al. Inferior epigastric artery pseudoaneurysm: A complication of paracentesis. *J Vasc Surg* 1998;28:566-9.
8. Todd AW. Inadvertent puncture of inferior epigastric artery during needle biopsy with fatal outcome. *Clin Radiol* 2001;56:989-90.
9. Satija B, Kumar S, Duggal RK, Kohli S. Deep circumflex iliac artery pseudoaneurysm as a complication of paracentesis. *J Clin Imaging Sci* 2012;2:10.

CASE REPORT

Lesch Nyhan Syndrome

Sridhar N*, Rambabu Sharma**, Ganesh Choudhary***, Suresh Dhaka****

INTRODUCTION

Lesch-Nyhan syndrome (LNS) is a rare inherited disorder caused by a deficiency of the enzyme hypoxanthine-guanine phosphoribosyl transferase-1 (HGPRT-1), produced by mutations in the HPRT gene located on the X-chromosome. LNS affect about one in 380,000 live births. The disease process mainly affects the male child and females are asymptomatic carriers. HGPRT deficiency leads to the characteristic triad of features: Hyperuricemia, Spectrum of neurological dysfunctions, and Cognitive and behavioural disturbances¹.

Self-injuring behaviour (SIB), it is rarely a presenting feature but eventually develops in nearly all cases. Affected patients are cognitively impaired and have behavioural disturbances that emerge between two and three years of age². Overproduction of uric acid “orange sand” in the diapers - uric acid crystalluria is due to HGPRT deficiency that causes a build-up of uric acid in all body fluids. Increased synthesis and decreased utilization of purines leads to high levels of uric acid production. This results in both high levels of uric acid in the blood and urine, associated with severe gout and kidney problems such as renal failure or frank hematuria-nephrolithiasis^{3,4}. Clinical manifestations according to age: At birth- no apparent neurologic dysfunction. After several months developmental retardation and neurologic signs become apparent. Before 4 months-hypotonia, recurrent vomiting and by age of 8-12 months-extra pyramidal signs prominent. The age at onset of self-injury may be as early as 1 year and occasionally as late as the teens.^{5,6}

Physical examination may show: growth

retardation, cognitive dysfunction and average IQ=60, all patients are wheelchair bound, self-mutilation like partial amputations of the fingers, lips, tongue, or oral mucosa, scarring from repetitive self-abrasion or hitting, in addition due to high levels of uric acid gouty arthritis and arthritic tophi may be seen. Lab. Studies can show hyperuricemia, hyperuricosuria, macrocytic anaemia. The diagnosis of Lesch-Nyhan syndrome may be confirmed by a thorough clinical evaluation, including a detailed patient history and specialized blood tests. The absence of the HPRT enzyme in cells from any tissue confirms the diagnosis. Molecular genetic testing for the HPRT1 gene is available to determine the specific disease-causing mutation. Carrier testing for Lesch-Nyhan syndrome is possible using molecular genetic testing^{5,6}. Prenatal diagnosis and preimplantation genetic diagnosis are possible if the disease causing HPRT1 gene mutation has been identified in an affected family member. Prenatal diagnosis can also be done by enzyme analysis. We report a case of LNS in a 22months old child who presented with buccal mucosal ulcers secondary to self-bites.

CASE REPORT

A 22 months old boy presented to our institution with history of motor development delay, dystonia and excessive irritability since 4 months of age. Self biting of fingers since 2 months, buccal mucosal ulcers since 20 days. He was the third child of healthy nonconsanguineous parents. There was no history of similar illness in first- degree and second-degree family members. The prenatal and neonatal periods were uneventful. At admission his length, weight and head circumference were 83 cm (–1 SD), 9.2 kg (–2 SD), and

*Resident, Department of Paediatric Medicine, SMS Medical College, Jaipur.

**Senior Professor, Department of Paediatric Medicine, SMS Medical College, Jaipur.

***Junior Specialist, Department of Paediatrics, Pandit Deendayal Upadhyaya Hospital, SMS Medical College Jaipur

****Senior Resident Department of Paediatric Medicine, SMS Medical College, Jaipur.

Corresponding Author

Dr. Ganesh Choudhary, Department of Paediatrics, Pandit Deendayal Upadhyaya Hospital, SMS Medical College Jaipur.

Mob. 9983467778

Mail: drganeshchoudhary@yahoo.com

48 cm (median) respectively. By 6 months of age, his head control was poor and he had abnormal dystonic movements and extensor spasms with marked hypotonia. Laboratory investigations including hemogram, thyroid function tests, liver function tests, computed tomography scan of cervical region were within normal limits, Brain magnetic resonance imaging (MRI) showed no signs of cortical atrophy or abnormal intensities except for arachnoid cyst, MR Spectroscopy was found to be normal. The patient developed hyperuricemia (uric acid [UA], 10.1 mg/dL) and increased urinary UA/creatinine ratio, 4.4 [control range 2.0]. EEG-Normal, Ammonia-24 micro mol/L (Control range-54, Lactate-13.0 mg/dl (Control range 4.5-19.8)

Child was treated with physiotherapy. He attained roll over at 14 months of age and started crying for his daily needs. But he started biting his own fingers at 19 months of age with developmental delay, we suspected the possibility of LNS. Laboratory investigations showed hyperuricemia, increased urine uric acid to creatinine ratio. Finally the diagnosis was confirmed by clinical exome sequencing. The report shows hemizygous nonsense variation in exon 7 of HPRT1 gene (ChrX:g.134498412134498412C>T;Depth:82x) that results in a stop codon premature truncation of protein at codon 170 (p.Arg170Ter;ENST00000298556.8) was detected. To know whether it's complete or incomplete HPRT deficiency levels to be determined. No such facilities are available in India. Hence we are planning to send the sample to United Kingdom.

On follow-up child is showing remarkable improvement in self-injurious behavior. Rest of clinical picture remains same. Beside the pharmacotherapy child is also receiving physiotherapy. Child was referred to dentist and he prescribed him tooth guard for protection of tongue, gums and teeth.

DISCUSSION

LNS is a rare disorder but it can easily be diagnosed by investigations like serum uric acid and urine uric acid to creatinine ratios available at most of the diagnostic centres.

Diagnosis of LNS is based on HPRT enzyme activity, preferably measured in live cells such as cultured fibroblasts, and on molecular genetic techniques demonstrating the gene mutation. Results might provide predictive clues about ultimate disease severity in

addition clinical and biochemical (hyperuricemia and hyperuricosuria), together with psychomotor signs of HPRT deficiency. An orange crystal in the diapers of a newborn is one of the early clues of the disease that may be observed. Overproduction of uric acid may lead to the development of uric acid crystals or stones in the kidneys, ureters, or bladder. Such crystals deposited in joints later in the disease may produce gout-like arthritis, with swelling and tenderness.

Self-Injury Behaviour (SIB) in LNS can be differentiated from SIB associated with autism or developmental disabilities by its sudden and more severe onset. Allopurinol will lower uric acid levels to normal but does not affect the behavioral aspects of the disease. Treatment options for management of SIB are physical restraints, behavioral, and pharmacological treatment (benzodiazepines) but the success rate is limited⁷. Physical restraints have been the sole reliable resource for preventing SIB. Cloth body restraints, cloth mittens, and plastic arm splints have all been useful in reducing the frequency of injury. Drastic measures, such as removal of the teeth or provision of tooth guards, are often taken to prevent further tissue damage. However, in many cases even with physical restraints, self-injury continues. Our patient's self-injurious behavior controlled with both pharmacotherapy and physical restraints. The aggressive behaviour usually wanes in patients older than 10-12 years of age. Patients with LNS usually die in their late second or third decade. The cause of death is renal failure or infections that are a result of decrease in lymphocyte and immunoglobulin G levels. With optimal care, few patients live beyond 40 years and most are confined to a wheelchair⁸.

REFERENCES

1. Torres RJ, Puig JG, Jinnah HA. Update on the phenotypic spectrum of Lesch-Nyhan disease and its attenuated variants. *Curr Rheumatol Rep* 2012;14:189-94.
2. Anderson LT, Ernst M (1994) Self-injury in Lesch-Nyhan disease. *J Autism Dev Disord* 24: 67-81
3. Nyhan WL, O'Neill JL, Jinnah HA, Harris JC (2014) Lesch-Nyhan Syndrome. *Gene Reviews*.
4. Augoustides-Savvopoulou P, Papachristou F, Fairbanks LD, Dimitrakopoulos K, Marinaki AM, et al. (2002) Partial hypoxanthine-Guanine phosphoribosyl transferase deficiency as the

- unsuspected cause of renal disease spanning three generations: a cautionary tale. *Pediatrics* 109: E17.
5. Lesch M, Nyhan WL (1964) A familial disorder of uric acid metabolism and central nervous system function. *Am J Med* 36: 561-70.
 6. Jinnah HA, Ceballos-Picot I, Torres RJ, Visser JE, Schretlen DJ, et al. (2010) LeschNyhan Disease international study group. Attenuated variants of Lesch-Nyhan disease. *Brain* 133: 671-89.
 7. Anderson LT, Ernst M. Self-injury in Lesch-Nyhan disease. *J Autism Dev Disord* 1994;24:67-81.
 8. Kale A, Shah K, Hallikerimath S. Lesch-Nyhan syndrome: A case report. *J Indian SocPedodPrev Dent* 2008;26 Suppl 1:S11-3





The Rajasthan Medical Journal

Guidelines / Instructions

Articles are accepted on the basis of significance, scientific merit and applicability. Authors are requested to make their report on the basis of original research carried out by themselves or their groups. The manuscripts will be reviewed for possible publication with the understanding that they are being submitted only to The Rajasthan Medical Journal and have not been published, simultaneously submitted, or already accepted for publication elsewhere. The editors review all submitted manuscripts initially and reject outright manuscripts that do not carry a clear message to the intended readership or are of insufficient originality.

The RMJ follows a policy of blind, peer review without revealing the identity of the contributors and each manuscript typically is read by at least two or three reviewers with expertise in the study's subject matter and methodology. All comments by reviewers are confidential and shall not be published. Within a period of around twelve weeks, the contributors will be informed about the reviewer's comments and acceptance or rejection of the manuscript revisions, if any, suggested by the editors or reviewers are communicated to the author's online requesting submission of the revised manuscript. The editor will make every effort to insure that authors informed as to the disposition of their manuscript within three months of receipt of the submission. The RMJ reserves the right to copy-edit the articles for length, grammar, punctuation, print style and format. Manuscripts must confirm to the guidelines for avoiding ethnic biases and sexist language. Research involved by an Institutional review board.

Online Submission of Manuscripts

Authors should submit their manuscript to The RMJ through its website www.rmjonline.in using the link "Submit Article". Electronic submission substantially reduces the editorial processing and peer reviewing times and shortens overall publication time.

Note: If you encounter any difficulty in submitting the manuscripts through the website link, you can e-mail all the manuscripts to the Editor as an attachment at editor@rmjonline.in

For online submission, articles should be prepared in two files (First Page file and The Article file). Images should also be attached with these two files.

First Page File: Prepare the title page, covering letter, acknowledgement, etc. using a word processor program. All

information which can reveal your identity should be here. Use text/doc/rtf/doc/PDF files. Do not zip the files.

The Article File: The main text of the article beginning from abstract till references (including tables) should be in this file. Do not include any other information (acknowledgment, your name in page headers, etc.) in this file. Use text /rtf/doc/PDF files. Do not zip the files. Limit the file size to 400 KB. Do not incorporate images in the file. If the file size is large, graphs can be submitted as images separately without incorporating them in the article file so as to reduce the size of the file.

Images: Submit good quality color images please! Each image should be less than 100 KB in size. Size of the image can be reduced by decreasing the actual height and width of the image (keep up to 400 pixels or 3 inches). JPEG image format is the most acceptable. Do not zip the files.

Legends: For the figures/images should be kept ready for copy paste during the submission process. Once a manuscript is submitted online, please send hard copies of the image and Contributors' Form to the Editor's address within four weeks. A Compact Disk or a Pen Drive containing the manuscript should be avoided.

Instructions for authors regarding manuscript submission

- Before you submit your manuscript, please read the editorial policy and the editorial guidelines of The RMJ carefully and make sure that your manuscript confirms to them. Manuscripts may be returned to the authors if they violate The RMJ editorial policy or deviate from its editorial guidelines.
- In order to widen the scope of communication, the following formats are made available - Editorials, review articles, original papers, leading articles, post graduate clinical updates, general practitioner updates, clinico-pathological conferences, unusual cases - case reports, clinical puzzles, readers corner and so on.

Physical Submission of Manuscripts

Send three copies of the manuscript, three sets of figures along with a covering letter, Contributors Form duly signed by all the contributors, checklist and a compact disk. Place the figures in a separate envelope. The covering letter must include information on prior or duplicate publication or submission elsewhere of any part of the work/study; and a statement of financial or other relationships that might lead to a conflict of interest. Copies of any permission(s) to reproduce published material, and to use illustrations or report information about identifiable people must accompany the manuscript. Send all the material to:



The Rajasthan Medical Journal

The Editor,

The Rajasthan Medical Journal (RMJ),

S.M.S. Medical College,

J.L.N. Marg, Jaipur – 302 004, Rajasthan, INDIA

It is essential that you provide complete contact details including full institutional address, telephone numbers, complete postal address and e-mail address for the corresponding author.

Type of Manuscripts and Word Limits

Leading Articles: The editors commission leading articles that are 1,000-1,500 words in length and address topics of current interest. They should be accompanied by no more than 10 references. Submissions are subjected to editorial review before acceptance. The editor retains the right to alter style and shorten material for publication.

Review Articles: The editorial board encourages submission of review articles on topics of current interest. The manuscript should be restricted to 4,000 words and up to 50 references. An abstract of no more than 250 words and up to 6 keywords should be provided.

Original Papers: These should be in the format of a) Introduction, b) Patients and methods, c) Results, and d) Discussion. The manuscript should be restricted to 3,000 words, up to 30 references and carry no more than 4 tables or figures. Please provide a 250 words structured abstract and up to 6 keywords. Structured abstract should include - background, study objective, design, setting, sample/population, duration of study, results and conclusion. Below the abstract, provide up to 6 keywords that will assist cross-indexing the articles and may be published with the abstract.

Unusual Cases: This section presents reports on rare cases. The manuscript should be in the format of a) Introduction, b) Case report, and c) Discussion. Submissions to this section should carry no more than 1,500 words, 2 figures and 5 references. An unstructured abstract of up to 150 words and 6 keywords should be provided.

Reader's Corner: Comments on papers recently published in the Journal. The letters should be restricted to up to 500 words and 3 references and should not carry figures.

Post Graduate Clinical Update: This category is for a concise, comprehensive summary of a topic of interest for post graduate students. The word count should not exceed 4,000

words.

Clinical Puzzle: This category is for unusual cases that makes an education point. Since the aim of these articles is to stimulate the reader to think about the case, the title should be ambiguous and not give away the final diagnosis immediately. Clinical puzzle should be in two parts. The first part should contain a brief description of the case (maximum 250 words) followed by an image and question-designed as task 1, 2, 3, 4, and so no. The legend of the image should not indicate the diagnosis but should simply describe the nature of the image. Each task is to be followed by a short description of what actually was done in the specific case. The answers to all the tasks should come in the second part, followed by a brief key message indicating few learning points. Clinical puzzle will not include more than 5 references, a maximum of 2 images, 2 figures and an author limit of 5. The quality of the image should be no less than 600 dpi and should be in JPG format. We do ask that authors indicate that they have obtained patient consent.

General Practitioner Update: This category presents a simple concise and clear summary of a topic of interest to general practitioners. The word count should not exceed 4,000 words. The word count excludes the title page, abstract, tables, acknowledgements & contributors, and the references.

Preparation of the Manuscript

Send laser print-out on thick paper of A4 size (8.27" x 11.69"), with margins of 1 inch on all side. Print on only one side of the paper. Use double spacing throughout. Number the pages consecutively, beginning with the title page. The language should be American English.

Title Page: Should carry the following information:

- Title of the article
- Running title or short title of no more than 50 characters
- Name of each contributor (Last name, First Name and initial of Middle Name)
- Name of the department(s) and institution(s) to which the work should be attributed
- Name, address, phone number, facsimile number and e-mail address of the contributor responsible for correspondence
- Type of manuscript (Original/Review/Unusual Cases, etc.)
- Word counts separately for abstract and for the text



The Rajasthan Medical Journal

(excluding the references and abstract)

Acknowledgement: Specify contributions that need an acknowledgement but do not justify authorship, such as general support by a departmental head or acknowledgements of technical, financial and material support and if the manuscript was presented as a part at the meeting in the organization, place and exact date on which it was read.

Abstract Page: The second page should carry the full title of the manuscript and an abstract (see above for word limits). For original articles, the abstract should be structured and arranged in the format indicated earlier.

Text of the Article:

- State the purpose of the article and summarize the rationale for the study or observation in the introduction. For unusual cases, give incidence of similar cases in the past.
- Describe the selection of the observational or experimental subjects clearly in patients and methods section. Identify the age, sex and other important characteristics of the subject. Identify the methods, apparatus (give the manufacturer's name in parentheses) and procedures in sufficient detail.
- Give references to established method, describe new or substantially modified methods, give reasons for using them, and evaluate their limitation.
- Identify precisely all drugs and chemicals used, including generic name(s) dose(s) and route(s) of admission.
- Report of randomized clinical trials should be based on the consort statement.
- Do not use patients' names, initials or hospital numbers especially in the illustrative material.
- Present the results in logical sequences in the text, tables and illustrations; emphasize or summarize only important observations. Use standard guidelines for statistics.
- Emphasize the new and important aspects of the study and the conclusion that follow from them along with implications of the findings and their limitation in the discussion section.

References:

- References should be numbered consecutively in the order in which they are first mentioned in the text.
- Identify references in text, tables, and legends by Arabic

numbers in superscript. References cited only in tables or figure legends should be numbered in accordance with the sequence established by the first identification in the text of the particular table or figure.

- The titles of journals should be abbreviated according to the style used in Index Medicus.
- Avoid using abstracts, un-published observations and personal communications as references.

Tables:

- Table should be self-explanatory and should not duplicate textual material.
- Tables with more than 10 columns and 25 rows are not acceptable. Limit the number to minimum required. Number tables in Arabic numerals, consecutively in the order of their first citation in the text and supply a brief title for each.
- Place explanatory matter in footnotes, not in the heading. Explain in footnotes all non-standard abbreviations that are used in each table. For footnotes, use the following symbols in the sequence: *, #, \$, **.
- Obtain permission for all fully borrowed, adapted and modified tables and provide a credit line in the footnote.

Illustrations (Figures):

- Submit three sets of sharp, glossy, un-mounted, color photographic prints, with height of 4 inches and width of 6 inches. Computerized color print-outs are not acceptable.
- Figure should be numbered consecutively according to the order in which they have been first cited in the text.
- Each figure should have a label pasted on its back indicating the number of the figure, the running title, top of the figure and the legends of the figure. Do not write on the back of figures, scratch, or mark them by using paper clips.
- Symbols, arrows or letters used in photomicrographs should contrast with the background and should be marked neatly with the background and should be marked neatly with transfer type or by tissue overlay and not by pen.
- If a figure has been published, acknowledge the original source and submit written permission from the copyright holder to reproduce the material. A credit line should appear in the legend for such figures.
- Type or print out legends (maximum 40 words,



The Rajasthan Medical Journal

excluding the credit line) for illustrations with Arabic numerals corresponding to the illustrations. When symbols, arrows or letters are used to identify parts of the illustration, identify and explain each one clearly in the legend explain the internal scale and identify the method of staining in photomicrographs.

Submitting a Revised Manuscript

While submitting a revised manuscript, contributors are requested to include, along with a single copy of the final revised manuscript, a photocopy of the revised manuscript with the changes underlined in red and with the point to point clarification to each comment. The manuscript number should be written on each of these documents.

Reprints

Journal does not provide any free printed reprints. Reprints can be purchased at the time of submitting the proofs.

Checklist (To be tick marked as applicable and one copy attached with the manuscript)

- Covering letter duly signed by all the contributors
- Previous publications / presentations mentioned
- Source of funding mentioned
- Conflicts of interest disclosed
- Authors
- a. Middle Name initials provided
- b. Author for correspondence, with e-mail address provided
- c. Identity not revealed in paper except title page (e.g. Name of the institute in material and methods, citing previous study as 'our study', name of institute in photographs, etc.)
- Presentation and format
- a. Double spacing

- b. Margins 2.5 cm from all four sides
- c. Title page contains all the desired information (vide supra)
- d. Running title provided (not more than 50 characters)
- e. Abstract page contains the full title of the manuscript
- f. Abstract provided (150 words for case reports and 250 words for original articles)
- g. Structured abstract provided for an original article
- h. Keywords provided (three or more)
- i. Headings in title case (not ALL CAPITALS, not underlined)
- j. Reference cited in superscript in the text without brackets
- k. Reference according to the journal's instructions
 - Language and Grammar
 - a. Uniformly American English
 - b. Abbreviations spelt out in full for the first time
 - Tables and Figures
 - a. No repetition of data in tables/graphs and in the text
 - b. Actual numbers from which graphs drawn to be provided
 - c. Figures necessary and of good quality (color)
 - d. Labels pasted on back of the figures (no names written)
 - e. Figure legends provided (not more than 40 words)
 - f. Patients' privacy maintained (if not, written permission enclosed)
 - g. Table and figure numbers in Arabic letter (not Roman)
 - h. Credit note for borrowed figures / tables provided
 - Manuscript provided on a Compact Disk



The Rajasthan Medical Journal

Contributors Form (to be modified as applicable and one single copy attached with the manuscript)

Title: _____

Manuscript Submission date _____ (for manuscripts submitted online) I / we certify that I / we have participated sufficiently in the intellectual content, conception and design of this work or the analysis and interpretation of the data (when applicable), as well as the writing of the manuscript, to take public responsibility for it and have agreed to have my / our name listed as a contributor. I / we believe the manuscript presents valid work. Neither this manuscript nor one substantially similar content under my / our authorship has been published or is being considered for publication elsewhere, except as described in the covering letter. I / we certify that all the data collated during the study is presented in this manuscript and no data from the study has been or will be published separately. I / we attest that, if requested by the editors. I / we will provide the data // information or will cooperate fully in obtaining and providing the data/information on which the manuscript is based, for examination by the editors or their assignees. I / we also certify that we have taken all necessary permissions from our institution and / or department for conduction and publishing the present work. Financial interests, direct or indirect, that exist or may be perceived to exist for individual contributors in connection with the content of this paper have been disclosed in the cover letter. Sources of outside support of the project are named in the cover letter.

I / we hereby transfer (s), assign(s), or otherwise convey(s) all copyright ownership, including and all rights incidental there to, exclusively to The Rajasthan Medical Journal (The RMJ) in the event that such work is published by The RMJ. The RMJ shall own the work, including 1) copyright; 2) the right to grant permission to republish the article in whole or in part, with or without fee; 3) the right to produce preprints or reprints and translate into languages other than English for sale or free distribution; and 4) the right to republish the work in a collection of articles in any other mechanical or electronic format. We give the rights to the corresponding author to make necessary changes as per the request of the journal, do the rest of the correspondence on our behalf and he/she act as the guarantor for the manuscript on our behalf. All persons who have made substantial contributions to the work reported in the manuscript, but who are not contributors, are named in the acknowledgment and have given me/us their written permission to be named. If I/we do not included an acknowledgment that names I / we have not received substantial contributions from non-contributors and no contributor has been omitted.

	Name	Signature	Date Signed
1	_____	_____	_____
2	_____	_____	_____
3	_____	_____	_____
4	_____	_____	_____
5	_____	_____	_____
6	_____	_____	_____



The Rajasthan Medical Journal

N. B. If you have already renewed your subscription for 2020, please disregard this notice

Editor

The Rajasthan Medical Journal

Name _____ Number of Copies _____

Organization (if applicable) _____ Department: _____

Position: _____ Designation _____

Address: _____

Contact No. _____ Fax _____

Email _____

Please circle the rate you are wanting

- | | |
|----------------------------|----------|
| 1. Annual Membership Fee - | Rs. 500 |
| 2. Life Membership Fee – | Rs. 2000 |

Thank you for renewing your details. The RMJ will send you an invoice for 2017

Please find enclosed cash / Cheque / DD No. _____ Dated _____

Drawn on (Bank Name) _____

For Rs. _____ favoring the Principal, SMS Medical College, Jaipur

Please invoice ☐

New Subscription ☐

Renewal ☐

